

## EFFECT OF LIQUID DISTILLERY DISCHARGES OF PEAK PRODUCTION SEASON ON SEED GERMINATION AND VIGOUR INDEX OF RARE PLANT *Withania somnifera* Dunal

N.K. MISHRA<sup>a1</sup> AND R.K. GUPTA<sup>b</sup>

<sup>a</sup>Department of Botany, A.B.R. P.G.College, Anapara, Sonebhadra, U.P., India  
E-mail : neelkanth\_2007@rediffmail.com

<sup>b</sup>Department of Botany, Postgraduate College, Ghazipur, U.P., India  
E-mail: drkpggc@gmail.com

### ABSTRACT

Present paper deals with the physiological response of distillery effluent on a rare medicinal plant *Withania somnifera* Dunal (Solanaceae) in connection with its conservation. This plant is at the verge of extinction due to its over-exploitation to fulfil the demands of pharmaceutical industries because of its higher medicinal values. This work was undertaken to study the effect of distillery effluent on two parameters of growth namely, seed germination and vigour index. The observations showed that in average concentration of the distillery effluent ranging between 25-30 % v/v, maximum vigour index was recorded whereas maximum seed germination was found in 5.0 % v/v strength of effluent. The vigour index was calculated by multiplying percentile germination and length of seedling. This study revealed that use of distillery effluent concentrations ranging from lower (5.0 % v/v) to average (25-30 % v/v) may be useful for conserving this plant.

**KEYWORDS:** Distillery discharges, vigour index, *Withania somnifera* Dunal

Rapid industrialization for sustaining economic stability is leading to pollute different factors of environment due to disposal of the untreated effluents in it however to ease the pressure and threatening level of industrial effluents on environment they are now being commonly used in agriculture practices for irrigation in dilute state. Somashekhar et al., (1998) put their view on this contradictory statement that the effluents may be beneficial or harmful to the crops, and it depends on type of industry from which the discharged effluents are being poured.

The effluents rich in organic and nitrogenous wastes may be profitable to the crops in preferable dilutions. Effluents from food industries, sugar mills and distilleries belong to beneficial group of industrial effluents. Another type of effluent is that which has higher concentration of chemicals, heavy metals and several other components fatal to life. This type of effluents are generally toxic to crop plants when poured into environment from chemical industries, tanneries, fertilizer factories etc.

In this communication, the effects of distillery effluent on growth of a well known medicinal plant *Withania somnifera* Dunal has been essayed. Mishra and Gupta, (2005) reported *Withania somnifera* Dunal to be in highly endangered state in surroundings of Ghazipur (U.P.). This species has become endangered due to over-exploitation and over-harvesting (Hemadri and Rao, 1998). For this sake, germination of seeds and vigour index of plantlets were taken into consideration.

Siddiqui and Goodwin, (2006) have contributed their works related to Vigour of different plant species. Gautam et al., (1992); Sahai and Srivastav, (1988); Sinha et al., (1988), Barman et al., (2001); Jayabalan et al., (2001) and Sprag et al., (2005) have evaluated the effects of different industrial effluents on various physiological aspects of plants.

### MATERIALS AND METHODS

Crude distillery effluent was collected from the outlet of Lord's Distillery Ltd., Nand Ganj, Ghazipur (U.P.) and was analysed for its physico-chemical properties following A.P.H.A. (1984). Table -1 depicts the report of this analysis. To determine effect of the distillery effluent it was diluted in de-ionized double distilled to prepare concentration grades ranging from 5.00% v/v, 10.00 % v/v, 15.00 % v/v to absolute 100.00 % v/v concentration.

The soil used in this test was somewhat modified common field soil which was thoroughly homogenized by light grinding and was mixed with fine sand in 3:1 ratio to prevent the problem of setting and layering of heavy and suspended solids of the effluent on the soil surface, then the soil was heated thrice in the oven at 120°C for 5 minutes to disinfect it properly against fungal, microbial and nematodal spores.

Seeds of *Withania somnifera* Dunal were collected from Botanical garden of institute of author's affiliation and were tested for their viability and germination in normal

---

<sup>1</sup>Corresponding author

conditions. Further, the seeds were presoaked overnight in different strengths of effluent and then thoroughly washed with distilled water, 50.00 % ethyl alcohol and again with distilled water for surface sterilization. Seeds were sown in twenty Petri-plates (marked for different concentration of effluent from 5.00-100.00 % v/v strength by which they were to be irrigated). A control was maintained for comparison and was irrigated with distilled water. Five replicas of this setup were taken for observation.

Emergence of radicle was taken as the criteria of germination. Germination pattern was recorded by 30 days daily inspection and the percentage of germination in each Petri-plate of every setup fortnightly by simple counting. The observations on length of seedling was done on a centimeter/ millimeter scale at stage just before the appearance of foliar leaves in the seedling. Vigour index was calculated.

## OBSERVATION

The percentile germination of *Withania somnifera* Dunal seeds was observed to be 60% in normal conditions (R.T. 30°C; Soaked in tap water) in control the germination started on 10th day with 20% germination at temperature ranging between 36°C to 44°C and was 60% on the eleventh day and it remained constant till the termination of experiment.

On the treatment with effluent, the percentile germination was maximum in 5.00% v/v strength of effluent, started on 6th day after sowing at temperature ranging between 34°C to 40°C with 10% germination which elevated to 60% on 15th day at temperature ranging between 37°C to 46°C. The germination was found to be lesser in higher concentration. A gradual decrease in germination percentage was noted with increase in strength of effluent. A drastic fall in germination percentage was noted in 35% v/v

**Table 1: Physico-chemicals properties of treated distilleries effluent**

S.N.	Property	Records	Remark
1.	Colour	Dark yellowish Brown	
2.	Odour	Sweet Foul	
3.	Temperature	34°C	
4.	Boiling Point	105°C	
5.	Freezing Point	- 4°C	
6.	pH	4.44	
7.	Conductivity	88.1768	
8.	Total solid	7.0092 g	per 25 ml.
9.	Total Dissolved solid	2.8910 g	per 25 ml
10.	Total suspended solid	4.1182 g	per 25 ml
11.	Oil & Grease	1.3260 g	per 25 ml
12.	BOD	30,800	
13.	Total Alkalinity	Nil (Acidic)	pH 4.44
14.	Dissolved Oxygen	Nil	
15.	Reducing Sugar	11.80 mg	per 25ml of 1% strength
16.	Non-reducing Sugar	20.45 mg	-do-
17.	Total Sugar	32.25 mg	-do-
18.	Bicarbonate	5.86 g	per 1000 ml
19.	Calcium	4.22 g	per 1000 ml
20.	Chloride	3.95 g	per 1000 ml
21.	Nitrate	37.00 mg	per 1000 ml
22.	Phosphate	22.00 mg	per 1000 ml

strength. The strength of effluent above 50.00% v/v was hazardous to the plant and strength above this range inhibited germination completely. (Table-2).

The observations on mean length of seedlings reflect that in control conditions it was observed to be 1.80 cM. The maximum mean length of seedlings was recorded in the strength of 25-30.00% v/v that is 2.80 cM. after which the harder strengths above 35.00% v/v cause unhealthy and stunted stature plantlets. The strength in range of 20.00-35.00% v/v was observed to be recommendable for getting good seedlings.(Table- 2).

Observations on the effects of distillery effluent on seedling germination and vigour index of *Withania somnifera* reveal that the diluted effluent (10.00% v/v 30.00% v/v strength) may be helpful for better cultivation, luxuriant growth and massive production of the plants of *Withania somnifera* because the value of vigour index is significantly elevated in this range. Maximum value (151.62) was found on 20.00% v/v strength. (Table-2).

## RESULTS AND DISCUSSION

Observations made on the effects of distillery effluent on seedling germination and vigour index of *Withania somnifera*, revealed that the diluted effluent (10.00-30.00% v/v strength) may be helpful for better cultivation, luxuriant growth and massive production of the

plants of *Withania somnifera* because the vigour index is higher (113.00-131.00) in this range which leads further to a healthy, luxuriantly growing plant while the germination was super fast and maximum in 5.00% v/v strength of effluent. The plantlets growing in 25.00% v/v and 30.00% v/v strength turned into healthy and vigorous plants because of significantly higher seedlings (2.80 cM.) and higher value of vigour index (151.62), which may establish pace between severe exploitation due to higher demands and production rate because of attaining the ability to produce more and more.

*Withania* faces the threat of extinction due to over-exploitation and relatively lesser production. This is how? The species may be escaped from the threat of endangerment and extinction. The higher concentrations above 50.00% v/v and crude discharges of distilleries may be a serious cause of the extinction of *Withania somnifera* like other plants.

## REFERENCES

- APHA AWWA WPCF, 1984. Standard Methods for the examination of water and waste water. 16th American Public Health Association, New York.
- Barman S.C., Kisku G.C., Salve P.R., Mishra D., Sahu R.K., Ramteke P.W. and Bhargava S.K. ,2001. Assessment of industrial effluent and its impact on

**Table 2 : Effect of different grades of concentration on germination of seeds, seedling growth and vigour index of *Withania somnifera* seeds**

S.N.	[A] Concentration Grade of Efflnt. (% v/v)	[B] Percentile Germination (%)	[C] Mean length of seedling (cM)	[D] Vigour index {[B]X[C]}
1.	Control	60	1.80	108.00
2.	05.0	60	1.83	109.80
3.	10.0	59	1.92	113.28
4.	15.0	57	1.99	113.43
5.	20.0	57	2.66	151.62
6.	25.0	54	2.80	151.20
7.	30.0	47	2.80	131.60
8.	35.0	30	2.60	78.00
9.	40.0	23	1.73	39.79
10.	45.0	14	1.20	16.80
11.	50.0	08	1.00	08.00
12.	55.0	00	0.00	00.00
13.	60.0	00	0.00	00.00

- soil and plants. J. Environ. Biol., **22**(4):251-256.
- Gautam D.D., Kumar K. and Beshnoi S. ,1992. Effect of dairy effluent on seed germination of some rohi and kharif crop plants. Jour. Environ. Biol., **13**(1): 7-12.
- Hemadri K. and Rao S.S. ,1998. Quantitative assessment of Medicinal plants II, Chittoor District, Andhra Pradesh, Manphar.
- Jayabalan M., Kannan D., Padmavathi S., Prem Kuma, G and Rajarathinam K. ,2001. Effect of tanary effluent on vigour index percentage seed germination index of seed of paddy ADT-36. J. Indl. Polln. Contl., **17**(2):207-212.
- Mishra N.K. and Gupta R.K. ,2005. Studies on Exploration and Conservation of Some Endangered Medicinal Plants Growing at Ghazipur (U.P.). XIV Annual Conference of PAS. Feb. 19-20, 2005, T.D. P.G. College, Jaunpur, U.P.India.
- Sahai R. and Srivastav C., 1988. Effect of fertilizer factory waste on germination , seedling growth and pigment content of plants. J. Indian Bot. Soc., **67** (3&4): 309-311.
- Siddiqui Md. Abdus and Goodwin P.B., 2006. Seed vigour in bean (*Phaseolus vulgaris* L. cv. Appolo) as influenced by Temperature and Water regime during Development and Maturation. Journal of Experimental Botany,**31** (1):313-323.
- Sinha B.K., Singh J.K. and Pandey A.K., 1988. Effect of distillery effluent supply on yield, tissue ascorbic acid, chlorophylls, catalase and peroxidase activities of sunflower (*Helianthus annus* L. vars. Indian and American) plants raised under standard soil pot culture conditions. Res. J. Pl. Environ., **4**(2):93-97.
- Somashekhar P.K., Gowda M.T.G., Shettigar S.L.N. and Srinath K.P. ,1988. Effect of Industrial effluents on crop plants. Ind. J. of Environ. Health, **26**(2) : 136-146.
- Sprag S.G., Kulkarni M.G., Light M.E., Van Staden J., 2005. Improving seedling vigour of indigenous medicinal plants with smoke. Bioresour Technol., **96**(12): 1323-1330.